

Electrocardiogram Testing in Symptomatic Patient with Type-II Diabetes and Left Ventricular Diastolic Dysfunction

ALI ZULQAR NAIN¹, AMBREEN SEHAR¹, MUSSARAT FATIMA²

¹MO at DHQ Hospital Sheikhupura.

²WMO at Jinnah Hospital, Lahore.

Corresponding author: Ali Zulqar Nain, Email: alizulqarnain89@gmail.com

ABSTRACT

Objective: The main objective of the study is to find the exercise electrocardiogram testing in asymptomatic patient with Type-II diabetes and left ventricular diastolic dysfunction.

Methods: This descriptive study was conducted in DHQ Hospital, Sheikhupura during January 2019 to November 2019. The data was collected from 50 patients of type II DM. These all patients has normal electrocardiogram (ECG) and normal systolic function.

Result: The data was collected from 50 patients of both genders. The mean age of the patients was 47.83 ± 5.61 years. After getting data the results were divided into two parts. There were 25 patients in group A and 25 patients group B. The mean duration of diabetes is 2.5 ± 5.61 years. There were 20 males and 30 females in this data. There were no significant difference of fasting blood sugar in both groups. There was no statically significant difference between two groups regarding left atrial dimension, aortic root dimension, LV end-diastolic dimension, ejection and LV mass.

Conclusion: It is concluded that It is concluded that coronary supply route sicknesses is a typical issue in DM, with diastolic brokenness and increment uniquely in patients with positive pressure practice electrocardiography.

Keywords: Exercise, Ventricular, Function

INTRODUCTION

Patients with type 2 diabetes often complain of fatigue and reduced exercise capacity which might be related to other illness conditions, for example, hypertensive left ventricular hypertrophy as well as coronary supply route sickness, and the resulting improvement of cardiovascular breakdown, however the presence of diabetes may freely add to the hindered practice limit. A few examinations have uncovered that activity limit diminishes in patients with type 2 diabetes and is related with cardiovascular mortality [1]. The proof demonstrates that in patients with type 2 diabetes, myocardial harm with fibrosis and apoptosis, disappointment in energy utilization, little vessel illness and autonomic neuropathies influence diastolic capacity before systolic capacity which is autonomous of hypertension and coronary corridor infection and prompts diabetic cardiomyopathy [2].

Individuals with diabetes (PWD) are at expanded danger of creating both miniature and macrovascular difficulties, which are lessened with legitimate glycaemic treatment. Contrasted and individuals without diabetes, men and, particularly, ladies with diabetes have diminished future (six to eight years less) [3]. One necessities to remember however that these individuals don't bite the dust from diabetes in essence but instead from cardiovascular sickness (CVD). At the hour of determination of type 2 diabetes (T2D), numerous patients as of now have at least one extra danger factors for macrovascular illness (stoutness, hypertension, dyslipidaemia, and smoking) and many have proof of plain atherosclerosis (past myocardial dead tissue (MI), ischaemic stroke, ischaemic changes on electrocardiogram (ECG), or fringe vascular infection) [4].

Doppler echocardiography is one of the most valuable clinical apparatuses for the appraisal of LV diastolic capacity. Doppler lists of LV filling are utilized for analytic purposes as well as for building up guess and assessing the impact of treatment. LV diastolic brokenness may speak to the primary phase of diabetic cardiomyopathy strengthening the significance of the early assessment of diastolic capacity in individual with diabetes [5].

Diabetes is related with expanded cardiovascular confusions, the most widely recognized of which are ischemic cardiomyopathy and LV brokenness. Diabetes is likewise connected with HF, basically through its association with hypertension and coronary vein disease.[8] Diabetes mellitus (DM) speeds up the cycle of coronary atherosclerosis and practical and basic heart disabilities [6].

DM isn't just a huge autonomous danger factor for the improvement of atherosclerotic ischemic coronary illness or

ventricular hypertrophy, yet it is likewise ready to trigger a diabetic cardiomyopathy because of some metabolic cycles: restraint of exchanging inside the cardiomyocyte from free unsaturated fat (FFA) to glucose digestion, dysregulation of FFA digestion with expanded take-up, diminished FFA oxidation, decrease of peroxisome proliferator-actuated receptor (PPAR), increment of PPAR-g and insulin-obstruction, and expanded intracellular lipogenesis which prompts cardiomyocyte lipotoxicity [7].

The main objective of the study is:

- To find the exercise electrocardiogram testing in asymptomatic patient with Type-II diabetes and left ventricular diastolic dysfunction.

MATERIALS AND METHODS

This descriptive study was conducted in DHQ hospital, Sheikhupura during January 2019 to November 2019. The data was collected from 50 patients of type II DM. These all patients has normal electrocardiogram (ECG) and normal systolic function.

Exclusion criteria

- Patients suffering from hypertension, ECG abnormalities and muscular disorder were excluded from this study.

Inclusion criteria

- All the patients who have type II DM, and ready to participate in the study.

Data collection: The data was collected from 50 patients. All patients went through complete reverberation Doppler study utilizing an industrially accessible echocardiography machine. LV measurements, left atrial measurements, divider thickness and launch part were estimated. Besides, transmitral stream was finished by beat wave Doppler between the tips of mitral valve flyers from four-chamber see measure top E wave speed, top A wave speed, E/A proportion, E wave deceleration time, and isovolumic unwinding time (IVRT). Exercise ECG was done to all patients to identify ischemic coronary illness and those with positive exercise ECG will inclined to coronary angiography.

The data was collected and analysed by using excel 2010.

RESULTS

The data was collected from 50 patients of both genders. The mean age of the patients was 47.83 ± 5.61 years. After getting data the results were divided into two parts. There were 25 patients in group A and 25 patients group B.

Group A: diabetic patients with LV diastolic dysfunction with negative stress ECG

Group B: patients with LV diastolic dysfunction with positive

stress ECG.

The mean duration of diabetes is 2.5±5.61 years. There were 20 males and 30 females in this data. There were no significant difference of fasting blood sugar in both groups. There was no statically huge contrast between two gatherings with respect to left atrial measurement, aortic root measurement, LV end-diastolic measurement, launch and LV mass. The exhibition of activity stress test has appeared in diabetics, a lower increment of E' speed (12,02±1,6 cm/sec versus 16,7±1,3 cm/s, p<0,01), slight increment of A' speed (p<0.01) contrasted and control subjects.

Table 1: Tissue Doppler Echocardiography characteristics of both groups after the stress test

Parameters	Group A	Group B	P
E' (cm/sec)	12.02±1,6	16.7±1.3	P<0.01
A' (cm/sec)	12.35±1,8	13.1±1.2	P<0.02
E'/A'	0.89±0.1	1.8±1.2	P<0.01
S' (cm/sec)	10.22±0.95	12.92±1.2	P<0.01

In group A there were 25 patients who were negative stress exercise in group A and 54% in group B. The statistical analysis showed a significant increase in the number of stress exercise ECG-positive patients among diabetic patients with diastolic dysfunction than diabetic patients without diastolic dysfunction (P<0.001).

Table 2: Stress exercise ECG in group A and group B

	Group A % age	Group B % age	χ ²	P-value
Exercise ECG			37	0.001
+ve	0	46		
-ve	100	54		
Functional			15.91	0.100
Capacity	0.1	9.1		
Poor	9.9	67.8		
Average	38.9	12.3		
Good	61.1	11.2		
Excellent				

DISCUSSION

Diabetes is notable to altogether expand CVD hazard, yet can't be viewed as a CHD same, because of extraordinary heterogeneity of the patients. All things considered, life-time danger of CHD is by all accounts very high in practically all individuals with the infection, which calls for individualized methodology and assessment for the presence and conceivable treatment of an incredible assortment of other as often as possible coinciding danger factors that can expand this danger [8]. It stays disrupted, on the grounds that all earlier randomized preliminaries have restrictions and are pointing towards equipoise, and accordingly, routine screening for quiet CHD in asymptomatic people with DM isn't right now suggested, as long as cardiovascular danger factors are dealt with [9]. Conversely, old style measures dependent on CE don't appear to have a similar capacity. Moreover, this irregularity is by all accounts related to the diabetic cardiomyopathy. Almost certainly, metabolic variations from the norm may assume a significant job [10]. Exploratory data from creature models of diabetes unequivocally uphold a causal part of insulin obstruction in the advancement of diastolic brokenness. Treatment with metformin forestalled the improvement of cardiomyocyte brokenness [11]. In an insulin-safe pre-diabetic rodent model, Mizushige et al. seen that the variations from the norm of diastolic filling happened before the advancement of straight to the point hyperglycaemia [12]. Histopathology contemplates proved expanded myocyte fibrosis and collagen statement, recommending that these auxiliary adjustments assume a significant function in the advancement of diastolic brokenness [13]. Another factor that may weaken diastolic

capacity is hyperglycaemia. There is trial proof that momentary hyperglycemia can modify cardiomyocyte contraction and unwinding. Moreover, high glucose focus causes the arrangement of

Aside from hazard factor treatment, be that as it may, the estimation of obtrusive treatment of coronary atherosclerosis (with the exception of the instance of intense coronary conditions) cutting edge glycation end products (AGE) that change collagen structure and meddle with intracellular calcium taking care of [14]. In people, Fang and partners found an expanded myocardial fibrosis in diabetic patients with heart brokenness. The current examination indicated that aortic sclerosis and mitral annular calcification were practically identical among various gatherings which mean that neither length of DM nor presence of diastolic brokenness has a function in their turn of events. Eren et al. expressed that aortic firmness is expanded in patients with hypertension, diabetes, or both even after the avoidance of coronary corridor sickness. Aortic firmness and LV diastolic brokenness are additionally related in these patients [15].

CONCLUSION

It is concluded that coronary supply route sicknesses is a typical issue in DM, with diastolic brokenness and increment uniquely in patients with positive pressure practice electrocardiography. Patients with positive pressure practice electrocardiography and coronary corridor infection apparent by coronary angiography had a practically identical diastolic capacity record to those with negative pressure practice electrocardiography. Doppler imaging application may legitimize routine screening for diastolic brokenness in diabetic patients dared to have healthy hearts.

REFERENCES

1. Vinereanu D, Nicolaides E, Tweddel AC., et al. Subclinical left ventricular dysfunction in asymptomatic patients with type II diabetes mellitus, related to serum lipids and glycated haemoglobin. Clin Sci 2003; 105: 591-599.
2. Diabetes Epidemiology Research International (DERI) Mortality Study Group: International evaluation of cause- specific mortality and IDDM. Diabetes Care. 1991; 14: 55- 60.
3. Mark KH, Faxon DP. Clinical studies on coronary revascularization in patients with type 2 diabetes. Eur Heart J. 2003; 24: 1087-1103.
4. Fang ZY, Yuda S, Anderson V., et al. Echocardiographic detection of early diabetic myocardial disease. J Am Coll Cardiol. 2003; 41: 611-617
5. Piccini JP, Klein L, Gheorghiade M, Bonow RO. New insights into diastolic heart failure: Role of diabetes mellitus. Am J Med 2004;116 Suppl 5A: 64S-75S
6. Poirier P, Bogaty P, Garneau C, Marois L, Dumesnil JG Diastolic dysfunction in normotensive men with well-controlled type 2 diabetes: Importance of maneuvers in echocardiographic screening for preclinical diabetic cardiomyopathy. Diabetes Care 2001;24:5-10.
7. Mytas DZ, Stogiannos PN, Zairis MN, Foussas SG, Pyrgakis VN, Kyriazis IA. Diabetic myocardial disease: Pathophysiology, early diagnosis and therapeutic options. J Diabetes Complications 2009;23:273-82
8. Nichols GA, Gullion CM, Koro CE, Ephross SA, Brown JB. The incidence of congestive heart failure in type 2 diabetes: An update. Diabetes Care 2004;27:1879-84
9. Bauters C, Lamblin N, Mc Fadden EP, Van Belle E, Millaire A, de Groote P. Influence of diabetes mellitus on heart failure risk and outcome. Cardiovasc Diabetol 2003;2:1.
10. Rathod MI, Sharma SK, Jain RK, Joshi RS, Mangudkar SS. A Study of Left Ventricular Diastolic Dysfunction in Patients with Diabetes Mellitus; 2010.
11. Iltercil A, Devereux RB, Roman MJ, Paranicas M, O'grady MJ, Welty TK, et al. Relationship of impaired glucose tolerance to left ventricular structure and function: The strong heart study. Am Heart J 2001;141:992-8.
12. Eren M, Gorgulu S, Uslu N, Celik S, Dagdeviren B, Tezel T. Relation between aortic stiffness and left ventricular diastolic function in patients with hypertension, diabetes, or both. Heart 2004;90:37-43
13. Grewal J, McCully RB, Kane GC, Lam C, Pellikka PA. Left ventricular function and exercise capacity. JAMA 2009;301:286-94
14. Kligfield P, Lauer MS. Exercise electrocardiogram testing: Beyond the ST segment. Circulation 2006;114:2070-82.
15. Mytas DZ, Stogiannos PN, Zairis MN, Foussas SG, Pyrgakis VN, Kyriazis IA. Diabetic myocardial disease: Pathophysiology, early diagnosis and therapeutic options. J Diabetes Complications 2009;23:273-82